



DISINFECTING WIPES

■ Specifications for Small Starter Kit (Item # 6185612)

- 2 each -180 wiper roll - 12" x 6" Spunlace Wipers
- 2 each -32 oz. bottle quaternary ammonium solution
- 2 each -1.3 gallon bucket with center slit lid to dispense wipers and resealable labels
- **Case Info** - 19.5" x 9.5" x 10" / Dimensional Weight: 10

■ Specifications for Large Starter Kit (Item # 6185613)

- 24 each -180 wiper roll - 12" x 6" Spunlace Wipers
- 24 each -32 oz. bottle quaternary ammonium solution
- 12 each -1.3 gallon bucket with center slit lid to dispense wipers and resealable labels
- **Case Info** - Weight is 200lbs and ships LTL on a pallet

■ Specifications for Refill Kit (Item # 6185614)

- 6 each -180 wiper roll - 12" x 6" Spunlace Wipers
- 6 each -32 oz. bottle quaternary ammonium solution
- 6 each -replacement resealable top label
- **Case Info** - 19" x 17" x 9" / Dimensional Weight: 19

Quaternary ammonium solution included with kit in accordance with CDC recommendations for disinfecting surfaces.

(Reference [CDC Fact Sheet for Disinfecting Your Facility](#)).



Freight: FOB Raleigh, NC Prepay and Add



DISINFECTING WIPES



Instructions

1. Place roll of wipes in bucket and pull center wipe through the center slit hole in the lid.
2. Pour 32 oz bottle of solution directly on the wipes in the bucket, and seal the bucket lid.
3. Cover the center slit hole with the resealable label while not using the wipes.

Active Ingredients

n-Alkyl (60% C14, 30% C16, 5% C12, 5% C18) dimethyl benzyl ammonium chlorides	0.105%	
n-Alkyl (68% C12, 32% C14) dimethyl ethylbenzyl ammonium chlorides	0.105%	
Inert Ingredients	99.790%	
TOTAL	100.000%	

Precautionary statements

HAZARDS TO HUMANS AND DOMESTIC ANIMALS

CAUTION: KEEP OUT OF REACH OF CHILDREN. Causes moderate eye irritation. Avoid contact with eyes or clothing. Wash thoroughly with soap and water after handling and before eating, drinking, chewing gum or using tobacco.

FIRST AID: Have the product container or label with you when calling a poison control center or doctor, or going for treatment.

If in eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes.

Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If swallowed: Call a poison control center or doctor immediately for treatment advice.

Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

If inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

NOTE TO PHYSICIAN

Probable mucosal damage may contraindicate the use of gastric lavage.

EPA REGISTERED:

EPA Reg. No. 1839-83-1124

EPA Est. No. 62511-KS-1